



Original Research Article

Studying the co-relation of preoperative serum albumin levels in emergency surgeries with rate of postoperative wound site infection

Nirmal Kumar Agarwal^{1,*}, Rajiv Mahato¹, Labanya Kumar Acharya¹, Subhash A Ragupathy¹

¹Dept. of General Surgery, Tezpur Medical College & Hospital, Tezpur, Assam, India



ARTICLE INFO

Article history:

Received 24-11-2022

Accepted 17-03-2022

Available online 31-07-2023

Keywords:

Emergency surgery

Serum albumin levels

Wound site infection

ABSTRACT

Aims: Hypoalbuminemia is a good marker of malnutrition. The objective is to analyze and correlate the rates of wound site infection in patient undergoing emergency surgery with preoperative serum albumin levels.

Materials and Methods: A prospective study of 98 patients who underwent emergency surgery between January 2021 to June 2021 in the Department of General Surgery, Tezpur medical college and hospital, Assam were considered for the study.

Serum albumin was determined preoperatively in all patients. Hypoalbuminemia was defined as serum albumin level of less than 3g/dl. Thirty day postoperative complication were analyzed. Wound site infection was confirmed by either by purulent discharge from the wound site or organism identified by aseptically obtained specimen from the superficial incision or from subcutaneous tissue by a culture based microbiological testing.

Results: A total of 98 patients who underwent emergency surgeries were studied. Out of 98 patients, 49 patients had normal Serum albumin levels and 49 patients had albumin levels of less than 3g/dl were included in the hypoalbuminemic group, Overall in the group of normal serum albumin level (n=49), patient with wound site infection were (n= 04) and patient without wound site infections were (n=45). In the group with level of albumin less than 3g/dl (n=49), patient with wound site infections (n=39), without wound site infections (n=10). After a chi-square test analysis with yates correction is 47.90.

Conclusions: Preoperative hypoalbuminemia is an independent risk factor and a predictor of postoperative wound site infections in patients undergoing emergency surgery. Identification and optimization of nutritional status prior to surgery may improve surgical outcomes

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

In a surgical patient, Malnutrition manifest most prominently as immunosuppression (an wound site infection and delayed wound healing).¹ Serum Albumin levels are useful in detecting and quantifying malnutrition.¹ Complication are defined as any deviation from the normal postoperative course.² It has been mentioned that

hypoalbuminemia is associated with higher rates of post-operative complication.³⁻⁵ Preoperative albumin levels of less than 3 g/dl is independently associated with an increased risk of developing serious complication within 30 days of surgery. Our Aim is to study the co-relation between preoperative albumin levels and post-operative wound site infection in patient undergoing emergency surgeries.

* Corresponding author.

E-mail address: nksurgeon@gmail.com (N. K. Agarwal).

2. Materials and Methods

This was a prospective study carried out from the period of January 2021–June 2021 in the Department of General Surgery in Tezpur Medical College and Hospital, Tezpur, Assam, A tertiary care hospital. All patients undergoing emergency surgeries were included in study and was studied for a postoperative period of 30 days. Patients less than 8 years of age, With hemoglobin less than 8g/dl and patients with Acute renal failure and People who lost follow up were excluded from study. Preoperative serum albumin was recorded for patient undergoing emergency surgeries.




Fig. 1: Work plan of the study

For the study patient were classified into two groups- Patients with normal serum albumin levels and Patients with low serum albumin levels in correlation with preoperative serum albumin levels (normal >3.0g/dl, Hypoalbuminemia <3.0g/dl). Patients demographic details, history, clinical findings, blood investigation and radiological investigations was done and were recorded. All patients were followed up or a period of 30 days. Patient were done regular dressing and was followed up for any complication.

The number of patients with wound site infection in both groups with hypoalbuminemia and normal albumin levels were studied using chi square test. All the data were analyzed using SPSS version 18. Wound site infection was confirmed by either purulent discharge from the wound site or organism identified by aseptically obtained specimen from the superficial incision or from subcutaneous tissue by a culture based microbiological testing.

3. Results

A total of 98 patients who underwent emergency surgeries were studied. Out of 98 patients, 49 patients had normal

Serum albumin levels and 49 patients had albumin levels of less than 3g/dl and were included in the hypoalbuminemic group.

Distribution of emergency surgeries done




Fig. 2: Distribution of emergency surgeries done

Overall in the group of normal serum albumin level (n=49), patient with wound site infection were (n= 04) and patient without wound site infections were (n=45). In the group with level of albumin less than 3g/dl (n=49), patient with wound site infections

(n=39) patient without wound site infections (n=10). The chi-square statistic with Yates correction is 47.9019. The

P value is < 0.00001. Significant at $p < .05$. The relationship between hypoalbuminemia and the wound site infection was found to be statistically significant. The wound site infection is more likely to occur when there is hypoalbuminemia than when serum albumin is normal.

4. Discussion

Complication After any surgery are inherent. Though it can be minimized it can never be eliminated. Many studies have shown that low preoperative serum albumin is a risk factor for postoperative complications.⁴⁻¹² In the present study we have established that Low serum albumin levels preoperatively is a very important risk actor for wound site infection in post operative period. When we compared the rates of wound site infection among the hypo-albuminemic group and individuals with normal serum albumin levels undergoing emergency surgery, the rates of wound site infection was statistically significant ($p < 0.05$). Low serum albumin is associated with poor wound healing, compromised collagen synthesis and a compromised immune response. The elements collectively may clarify the higher risk of wound site infection in hypoalbuminemic patients.

Concerning the treatment of preexisting low serum albumin levels few studies have shown no beneficial outcome with the administration of albumin infusion.^{13–15} This could be due to inadequate albumin replacement or increased leakage of synthetic albumin into extravascular spaces. However, recently some study highlighted that preoperative nutrition supplements may benefit the patients and improve surgical outcomes.^{16,17} We believe that this one of the precise series clearly showing an association between preoperative serum albumin and significant wound site infection after emergency surgery. Limitation of our study are; Single centered study design, small size study population, which could influence the ability to determine preoperative serum albumin as a predictor: Serum albumin which itself is a multifaceted variable which could be altered broadly in various physiopathological condition. Despite all the limitations, Preoperative serum albumin remained a significant predictor of post operative complications and mortality in patients undergoing emergency surgery. Hence to validate our findings further appropriately design studies are suggested.

5. Conclusion

Preoperative hypoalbuminemia is an independent risk factor for the development of low postoperative complication. Surgeons should be aware of the consequences of low preoperative serum albumin and concerns on nutritional optimisation which is likely to increase the postoperative outcome

6. Source of Funding

None.

7. Conflict of Interest

None.

8. Acknowledgement


The authors are grateful to all the patients, staffs, colleagues who assisted in the study.

References

1. Sullivan DH, Bopp MM, Roberson PK. Protein- energy undernutrition and life-threatening complications among the hospitalized elderly. *J Gen Intern Med.* 2002;17(12):923–32. doi:10.1046/j.1525-1497.2002.10930.
2. Clavien PA, Camargo CA, Croxford R, Langer B, Levy GA, Greig PD, et al. Definition and classification of negative outcomes in solid organ transplantation. Application in liver transplantation. *Ann Surg.* 1994;220(2):109–20. doi:10.1097/00000658-199408000-00002.
3. Lohsiriwat V, Lohsiriwat D, Boonnuch W, Chinswangwatanakul V, Akaraviputh T, Lert-Akayamane N, et al. Pre-operative hypoalbuminemia is a major risk factor for postoperative complications following rectal cancer surgery. *World J Gastroenterol.* 2008;14(8):1248–51. doi:10.3748/wjg.14.1248.
4. Kuzu MA, Terzioğlu H, Genç V, Özban M, Sonyürek P. Preoperative nutritional risk assessment in predicting postoperative outcome in


- patients undergoing major surgery. *World J Surg.* 2006;30(3):378–90. doi:10.1007/s00268005-0163-1.
5. Alves A, Panis Y, Mathieu P, Manton G, Kwiatkowski F, Slim K, et al. Postoperative mortality and morbidity in French patients undergoing colorectal surgery: results of a prospective multicenter study. *Arch Surg.* 2005;140(3):278–83.
6. Putwatanap R, Reodecha P, Sirapo-ngam Y, Lertsithichai P, Sumbonnanonda K. Nutrition screening tools and the prediction of postoperative infectious and wound complications: Comparison of methods in presence of risk adjustment. *Nutrition.* 2005;21(6):691–7. doi:10.1016/j.nut.2004.10.015.
7. Ryan AM, Hearty A, Prichard RS, Cunningham A, Rowley SP, Reynolds JV, et al. Association of hypoalbuminemia on the first postoperative day and complications following esophagectomy. *J Gastrointest Surg.* 2007;11(10):1355–60. doi:10.1007/s11605-007-0223-y.
8. Leite JF, Antunes CF, Monteiro JC, Pereira BT. Value of nutritional parameters in the prediction of postoperative complications in elective gastrointestinal surgery. *Br J Surg.* 1987;74(5):426–9. doi:10.1002/bjs.1800740538.
9. Neumayer L, Hosokawa P, Itani K, El-Tamer M, Henderson WG, Khuri SF, et al. Multivariable predictors of postoperative surgical site infection after general and vascular surgery: results from the patient safety in surgery study. *J Am Coll Surg.* 2007;204(6):1178–87. doi:10.1016/j.jamcollsurg.2007.03.022.
10. Pikul J, Sharpe MD, Lowndes R, Ghent CN. Degree of preoperative malnutrition is predictive of postoperative morbidity and mortality in liver transplant recipients. *Transplantation.* 1994;57(3):469–72. doi:10.1097/00007890-19940215000030.
11. Lai CC, You JF, Yeh CY, Chen JS, Tang R, Wang JY, et al. Low preoperative serum albumin in colon cancer: a risk factor for poor outcome. *Int J Colorectal Dis.* 2011;26(4):473–81. doi:10.1007/s00384-0101113-4.
12. Chai FY, Jiffre D. Preoperative hypoalbuminemia is an independent risk factor for the development of surgical site infection following gastrointestinal surgery. *Ann Surg.* 2011;254(4):665. doi:10.1097/SLA.0b013e31823062f3.
13. Cerantola Y, Grass F, Cristaudi A, Demartines N, Schäfer M, Hübner M, et al. Perioperative nutrition in abdominal surgery: recommendations and reality. *Gastroenterol Res Pract.* 2011;35(2):241–8. doi:10.1177/0148607110374477.
14. Bromenshenkel J, Newcomb M, Thompson J. Continuous quality improvement efforts decrease postoperative ileus rates. *J Healthc Qual Off Publ Natl Assoc Healthc Qual.* 2000;22(2):4–7.
15. Rubin H, Carlson S, Demeo M, Ganger D, Craig RM. Randomized, double-blind study of intravenous human albumin in hypoalbuminemic patients receiving total parenteral nutrition. *Crit Care Med.* 1997;25(2):249–52. doi:10.1097/00003246-199702000-00009.
16. Moskovitz DN, Kim YI. Does perioperative immunonutrition reduce postoperative complications in patients with gastrointestinal cancer undergoing operations? *Nutr Rev.* 2004;62(11):443–7. doi:10.1111/j.1753-4887.2004.tb00016.x.
17. Dempsey DT, Mullen JL, Buzby GP. The link between nutritional status and clinical outcome: can nutritional intervention modify it? *Am J Clin Nutr.* 1988;47(2):352–6.

Author biography

Nirmal Kumar Agarwal, Associate Professor  <https://orcid.org/0000-0001-7350-9754>

Rajiv Mahato, Assistant Professor  <https://orcid.org/0000-0003-2583-6050>

Labanya Kumar Acharya, Registrar  <https://orcid.org/0000-0002-0584-9228>

Subhash A Ragupathy, Post Graduate Trainee  <https://orcid.org/0000-0002-0253-1812>

Cite this article: Agarwal NK, Mahato R, Acharya LK, Ragupathy SA. Studying the co-relation of preoperative serum albumin levels in emergency surgeries with rate of postoperative wound site infection. *Panacea J Med Sci* 2023;13(2):295-298.