

## Sexual issues in dementia: An overview

Pragya Lodha<sup>1</sup>, Avinash De Sousa<sup>2\*</sup>

<sup>1</sup>Clinical Psychologist and Research Assistant, <sup>2</sup>Consultant Psychiatrist and Founder Trustee, Desousa Foundation, Mumbai, Maharashtra, India

\*Corresponding Author: Avinash De Sousa

Email: [avinashdes888@gmail.com](mailto:avinashdes888@gmail.com)

### Abstract

Dementias are the most common type of neurodegenerative disorders. Behavioral disturbances are seen in 50-80% of patients suffering from these disorders. Sexually inappropriate behaviors are not as common as other behaviors seen in dementia but may be seen in 7-30% subjects. These behaviors cause a lot of distress to all those who are affected along with shame and embarrassment. Sexual behaviors in dementia lie on a spectrum that may range from mild or benign to severe. There are no rigorous clinical studies for the treatment of these behaviors and there is no fixed classification for the types of behavior either. There is sparse data that suggest efficacy for some commonly used treatment modalities in dementia. In this chapter, we review the various aspects of these behaviors, their types and the available treatment for the same.

**Keywords:** Sexual issues, Dementia, Sexuality, Sexually inappropriate behavior.

### Introduction

Dementia is a neuropsychiatric disorder characterized by memory loss, impairments in cognition, behavioral disturbances and problems in activities of daily living. It is the most common disorder seen in geriatric patients affecting about 5-10% of people above the age of 65 years and 20-40% of people older than 75 years.<sup>1</sup> The most common form of dementia is Alzheimer's-type dementia (AD) followed by vascular dementia. Behavioral disturbances are common in dementia.<sup>2</sup> These can be defined as behaviors that are unsafe and disruptive and that interfere with the care of the patient in any given environment. Around 60-65% of patients with dementia will have behavioral disturbances at any one point in time.<sup>3</sup> These disturbances contribute to increased morbidity, greater health care resource utilization, and premature institutionalization.<sup>4</sup> The most distressing amongst the behavioral disturbances are sexual symptoms that can serve to be distressing as well as embarrassing for the family members.<sup>5</sup> This article aims to provide the busy clinician an overview of the various sexual symptoms that may be seen in dementia and methods employed to manage the same.

### Epidemiology

Studies have demonstrated that 7-30% of patients with dementia may exhibit sexual symptoms.<sup>6</sup> There are no large scale studies that are epidemiological in nature and that have been done to look at sexual symptoms per se. clinically though these symptoms may be commoner and are under-reported due to embarrassment and non-inquiry by clinicians that may result in an under-reporting of the same. Research in this regard is needed both in the Indian and western context.<sup>7</sup>

### Types of Sexually Inappropriate Behaviours in Dementia

There are various forms of inappropriate sexual behavior seen in dementia. These lie on a spectrum from mild to severe and may manifest in various forms.<sup>8-9</sup>

1. Sexual talk and foul language

2. Sexual jokes and sexually suggestive gestures and remarks.
3. Exposing oneself to others.
4. Public masturbation.
5. Kissing and hugging that exceeds what is normal.
6. Disrobing and moving about without clothes.
7. Touching the breasts, buttocks, and genitals of staff and family members.
8. Attempting sexual intercourse and oral sex.
9. Requesting genital care from the staff and family members.
10. Reading and watching pornographic material.
11. Sexually deviant behaviour.
12. Attempting to sexually assault staff and servants.
13. Collecting and hoarding sexual books and magazines.

### Causes of Sexually Inappropriate Behaviour in Dementia

#### Neurobiological Factors

Certain brain systems have been implicated in the neurobiology of inappropriate sexual behaviors in dementia viz. the frontal lobes, the temporo-limbic system, the striatum, and the hypothalamus. Each of these systems has specific roles in various sexually inappropriate behaviors that may be seen in dementia.<sup>10</sup>

#### Frontal Lobes

The frontal lobe has the sexual center of the brain and mediates the expression of sexual behaviors. Dysfunction of the frontal system typically involves disinhibition rather than hypersexuality which is a hallmark of sexually inappropriate behavior in dementia.<sup>11-12</sup>

#### Temporo-Limbic System

Sexual behaviors are mediated through the temporo-limbic system in both humans and animals. Both temporal lobes have been implicated in behaviors that are auto-erotic in nature and self-stimulatory behaviors that may be seen in dementia.<sup>13</sup> Hypersexual behaviors have also been reported after temporal lobe degeneration. The right temporal lobe has been implicated in altered sexual behaviors as it modulates

emotions and the understanding of the effect associated with sexual arousal.<sup>14</sup>

### **Striatum**

Sexual behaviors which are repetitive, habit like and obsessive in nature are associated with lesions of the striatum and its connections with the cerebral cortex.<sup>15</sup>

### **Hypothalamus**

Lesions to the hypothalamus can lead to an increase in sexual behavior. Increased sexual drive and hypersexuality is a hypothalamic dysfunction.<sup>16</sup> The right hypothalamus and periventricular area can cause hypersexuality as seen in mania.<sup>17</sup>

### **Psychological Factors**

Psychological factors for sexually inappropriate behavior in dementia are unclear. The expression of sexuality involves complex psychological, and environmental factors that may cause such behavior. Sexual manners are learned behaviors that may be forgotten as a result of dementia. In other cases, it may be related to a psychological need for intimacy that has been sexualized.<sup>18</sup> People with dementia may feel disconnected from others, and they may have lost the ability to speak or to communicate their desires and needs. Consequently, they may be acting out a strong need for human connection and touch which may be misinterpreted as sexual.<sup>19</sup>

Researchers have coined a term iatrogenic loneliness that has been induced by staff attitudes and organizational structures that discourage or fail to accommodate any form of intimate relationship within the institutional setting. This may be expressed as sexually inappropriate behavior by the patient with dementia.<sup>20</sup> The patient with dementia just craves the touch itself. People with dementia may confuse staff and other residents with a much-loved partner and respond out of that misinterpretation.<sup>21</sup>

### **Assessment of the Patient**

One must obtain a comprehensive history, including a thorough sexual history. History should be obtained from the caregivers or family members or patient if possible. It must be determined that these behaviors are truly sexual and inappropriate and not just a yearning for closeness and warmth. Misinterpretation by staff at nursing homes must be looked into. One must carry out a good mental status and physical examination.<sup>22</sup> Laboratory data including routine tests and neuroimaging studies may be done when needed. Neuropsychological testing may help determine the patient's level of cognitive functioning and extent of memory impairments.<sup>23</sup>

### **Management**

There are few systematic studies on the treatment of sexually inappropriate behavior in dementia.<sup>24</sup> Most of the data available to us are from anecdotal case reports or case series. The type of treatment depends upon the distress caused by the situation, type of sexually inappropriate behavior, underlying medical conditions and whether the patient is at home or institutionalized. Both non-pharmacological and

pharmacological treatments have been used in the management.<sup>25-26</sup>

### **Non Pharmacological Treatments**

One has to determine what social cues are misinterpreted and leads to behavior. Modification of these cues usually leads to a reduction in these inappropriate behaviors.

### **Supportive Psychotherapy**

Psychotherapy is especially useful for spouses of patients who have inappropriate behaviors. They often need reassurance and support that these behaviors are secondary to the illness and not a reflection of their relationship which may cause depression. It may also be useful to view their partner's sexual requests as a call for closeness and reassurance rather than just sexual.<sup>27</sup>

### **Behavior Modification**

When inappropriate behaviors occur, one must sensitively explain to the patient who can understand that such behaviors are unacceptable. One must prevent confrontation and should not cause excessive guilt or shame. Never ignore or laugh at the behavior as they may then be encouraged. Distraction may be a very useful technique in some cases.<sup>28-29</sup>

### **Environmental Changes**

In nursing homes, single rooms and provision for the spouse meeting the patient may help reduce the frequency of such behaviors. Avoidance of external cues such as over stimulating television or radio programs is helpful.<sup>30</sup> In patients with a tendency to expose themselves or masturbate in public, trousers that open in the back or that are without zippers may be helpful. Provision for adequate social activity is helpful.<sup>31</sup>

Changing the attitudes of the family, caregivers, and staff in nursing homes is very essential. The care of dementia patients at a nursing home demands a high degree of technical and interpersonal skills. Caregivers are often caught between moral norms, society, nursing home rules, a person's rights and providing appropriate care for their patients.<sup>32</sup>

Sex education programs for the family, the caregivers, and the staff at the nursing homes can add to the quality of life of a demented person. Emphasis on the need for normal sexual expression while preventing inappropriate sexual behaviors should be emphasized.<sup>33</sup>

### **Pharmacological Treatments**

There are no systematic studies or randomized double-blind placebo-controlled trials for any of the drugs that are used to treat sexually inappropriate behaviors in dementia.<sup>34</sup> Medication should only be used when all other treatment methods have failed. One must start medicine at a low dose and go slow in increasing the dose. One must monitor for side effects and inform staff and relatives about the same. Some medicines may precipitate or worsen these behaviors and must be discontinued or avoided. There are many medications that have been used in the management of such behaviors.<sup>35</sup>

### Selective Serotonin Reuptake Inhibitors

These medicines are known to decrease inappropriate behaviors by their effects on anxiety and obsessive thinking as well as their sexual side effects. They also reduce the levels of sex hormones and can treat comorbid depression and anxiety disorders. The common side effects seen with these medications are gastrointestinal disturbances, headache, insomnia, and sexual dysfunction. All the SSRIs have been used in small case series and anecdotal case reports in the management of these problems.<sup>36</sup> There have been case reports with Paroxetine showing some amount of success and Escitalopram too has been used.<sup>37-38</sup> There have been reports of successful elimination of such behaviors when Paroxetine has been combined with Clomipramine.<sup>39</sup> However side effects need monitoring and patients may need to be started on an antacid when on these medications

### Antipsychotics

There are no known clinical trials in the elderly on the use of antipsychotic medications in the treatment of these behaviors, but clinical experience and evidence point to their efficacy.<sup>40</sup> These drugs are thought to decrease sexually inappropriate behavior by their dopamine-blocking effects. Dopamine is a prosexual neurotransmitter. Atypical antipsychotics have been used far more than typical and have shown greater efficacy. Quetiapine and Risperidone have been used in most cases but side effects like extrapyramidal reactions and drowsiness must be monitored.<sup>41-42</sup> Also one has to be careful when using these drugs in patients with diabetes and hyperlipidemia. Risperidone has been used the maximum while case reports with Olanzapine and Amisulpride also exist.<sup>43-44</sup>

### Trazodone

Trazodone is an older antidepressant drug which is a presynaptic reuptake inhibitor and a mild postreceptor agonist of serotonin. Case series have reported efficacy of the drug in reducing sexually inappropriate behavior. The dose range for trazodone that may be used is 100 and 500 mg a day in divided doses. The response was thought to be due to the calming effect of the drug and not its antidepressant effect. One must monitor for side effects like headache, dry mouth, sedation, orthostatic hypotension, and weight gain while on the drug.<sup>45</sup> Priapism (painful erection) occurs in 1 in 6000 patients and can be distressing if it happens in a case of dementia or with an already existing prostate problem.<sup>46</sup>

### Anti-androgens

The commonly used antiandrogens are medroxyprogesterone acetate (MPA) and cyproterone acetate (CPA). They act by the reduction in serum testosterone level which will impair sexual functioning, and this, in turn, will eliminate the inappropriate behaviors. MPA is progesterone that decreases the level of testosterone by inhibiting the levels of pituitary luteinizing hormone (LH) and follicle stimulating hormone (FSH). The major side effects are sedation, increased appetite, weight gain, fatigue, loss of body hair, hot and cold flashes, mild diabetes, decreased ejaculatory volume, and

symptoms of depression. These drugs have sparse evidence in the form of case reports.<sup>47</sup>

### Estrogens

These medications act by reducing LH and FSH secretion and thereby reducing testosterone production. The common estrogens are diethylstilbestrol (DES) and conjugated estrogens. Common side effects include fluid retention, nausea, vomiting, impotence, and gynecomastia. One must be careful while starting any hormonal medications in older patients due to the various side effects present.<sup>48</sup>

### Gonadotrophin-releasing Hormone Analogs

These medications suppress testosterone production by stimulating the secretion of pituitary LH and FSH. This results in an increase in estrogen production, thereby decreasing the level of testosterone. Leuprolide acetate is the common gonadotrophin-releasing hormone (GnRH) analog used in clinical practice. These drugs must be used continuously to maintain their effectiveness.<sup>49</sup>

The use of hormonal agents for the treatment of inappropriate sexual behavior in the elderly is sensitive in many ways. The side-effect profile of these drugs and the social stigma associated with using them are seen as chemical castration. The same treatment has been used in sex offenders with high sex drives and thus must be used with caution in patients with dementia.<sup>50</sup>

### Cimetidine

Cimetidine is an H<sub>2</sub> receptor antagonist with antiandrogen effects. Anecdotal case reports exist in the management of patients with the drug though no major studies are available.<sup>51</sup>

### Pindolol

It is a beta blocker that has been used to augment SSRIs and in agitation and hypersexual behaviors. Common side effects of pindolol are fatigue and hypotension. This drug is thought to reduce inappropriate behavior by decreasing adrenergic drive and thus decreasing agitation, aggression, and inappropriate behavior.<sup>52</sup>

### Mood Stabilizers

There are no major reports on the use of mood stabilizers in the treatment of inappropriate sexual behaviors, though these are widely used in clinical practice. Common side effects of these medications are tremors, sedation, falls, and weight gain.<sup>53</sup>

### Cholinesterase Inhibitors

Cholinesterase inhibitors such as donepezil, rivastigmine, and galantamine have been found to be effective in treating cognitive dysfunction and behavioral disturbances associated with dementia. However, there are no reports on the use of these medications in the treatment of inappropriate sexual behaviors associated with dementia.<sup>54</sup>

### Memantine

The N-methyl D-aspartate (NMDA) receptor antagonist memantine has been approved for the use in moderate to severe dementia. There are no current reports on their use in elderly patients with inappropriate sexual behavior.<sup>55</sup>

### Conclusion

Dementia is a public health problem which shall escalate in the years to come. Behavioral problems associated with dementia are very common and are a major source of distress. These behaviors are also the most common reason for the placement of a patient suffering from dementia in nursing home care. There are limited data on the various aspects of these behaviors and their management. Future research should not only focus on effective treatment but also on early detection and prevention of such behavior and their sound pharmacological management. This will reduce undue suffering for both the patients and their caregivers and also enhance the quality of life and dementia care for all those concerned.

**Acknowledgments:** Nil.

**Conflict of Interest:** Nil.

**Funding:** Nil.

### References

1. Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W, Ferri CP. The global prevalence of dementia: a systematic review and meta-analysis. *Alz Dementia* 2013;9(1):63-75.
2. Chan KY, Wang W, Wu JJ, Liu L, Theodoratou E, Car J, et al. Epidemiology of Alzheimer's disease and other forms of dementia in China, 1990–2010: a systematic review and analysis. *Lancet* 2013;381(9882):2016-23.
3. Black W, Almeida OP. A systematic review of the association between the behavioral and psychological symptoms of dementia and the burden of care. *Int Psychogeriatr* 2004;16(3):295-315.
4. Azermi M, Petrovic M, Elseviers MM, Bourgeois J, Van Bortel LM, Vander Stichele RH. Systematic appraisal of dementia guidelines for the management of behavioral and psychological symptoms. *Ageing Res Rev* 2012;11(1):78-86.
5. Hillman J. Sexual issues and aging within the context of work with older adult patients. *Profess Psychol Res Pract* 2008;39(3):290-7.
6. Mahieu L, Gastmans C. Sexuality in institutionalized elderly persons: a systematic review of argument-based ethics literature. *Int Psychogeriatr* 2012;24(3):346-57.
7. Shaji KS, Jithu VP, Jyothi KS. Indian research on aging and dementia. *Indian J Psychiatry* 2010;52(Suppl1):S148-52.
8. Higgins A, Barker P, Begley CM. Hypersexuality and dementia: dealing with inappropriate sexual expression. *Br J Nurs* 2004;13(22):1330-4.
9. Johnson C, Knight C, Alderman N. Challenges associated with the definition and assessment of inappropriate sexual behavior amongst individuals with an acquired neurological impairment. *Brain Inj* 2006;20(7):687-93.
10. Tsatali MS, Tsolaki MN, Christodoulou TP, Papaliagkas VT. The complex nature of inappropriate sexual behaviors in patients with dementia: Can we put it into a frame?. *Sexuality Disabil* 2011;29(2):143-56.
11. Murtha S, Cismaru R, Waechter R, Chertkow H. Increased variability accompanies frontal lobe damage in dementia. *J Int Neuropsychol Soc* 2002;8(3):360-72.
12. Rosen HJ, Gorno-Tempini ML, Goldman WP, Perry RJ, Schuff N, Weiner M, et al. Patterns of brain atrophy in frontotemporal dementia and semantic dementia. *Neurol* 2002;58(2):198-208.
13. Chan D, Fox NC, Scahill RI, Crum WR, Whitwell JL, Leschziner G et al. Patterns of temporal lobe atrophy in semantic dementia and Alzheimer's disease. *Ann Neurol* 2001;49(4):433-42.
14. Bozeat S, Gregory CA, Ralph MA, Hodges JR. Which neuropsychiatric and behavioral features distinguish frontal and temporal variants of frontotemporal dementia from Alzheimer's disease?. *J Neurol Neurosurg Psychiatry* 2000;69(2):178-86.
15. Tsuboi Y, Uchikado H, Dickson DW. Neuropathology of Parkinson's disease dementia and dementia with Lewy bodies with reference to striatal pathology. *Parkinsons Related Disord* 2007;13: S221-4.
16. Piguet O, Petersén Å, Yin Ka Lam B, Gabery S, Murphy K, Hodges JR, et al. Eating and hypothalamus changes in behavioral-variant frontotemporal dementia. *Ann Neurol* 2011;69(2):312-9.
17. Ishunina TA, Kamphorst W, Swaab DF. Metabolic alterations in the hypothalamus and basal forebrain in vascular dementia. *J Neuropathol Experiment Neurol* 2004;63(12):1243-54.
18. Ward R, Vass AA, Aggarwal N, Garfield C, Cybyk B. A kiss is still a kiss? The construction of sexuality in dementia care. *Dementia* 2005;4(1):49-72.
19. Black B, Muralee S, Tampi RR. Inappropriate sexual behaviors in dementia. *J Geriatr Psychiatr Neurol* 2005;18(3):155-62.
20. Benbow SM, Beeston D. Sexuality, aging, and dementia. *Int Psychogeriatr* 2012;24(7):1026-33.
21. Series H, Degano P. Hypersexuality in dementia. *Adv Psychiatr Treat* 2005;11(6):424-31.
22. Wallace MA. Assessment of sexual health in older adults. *Am J Nurs* 2008;108(7):52-60.
23. Feldman HH, Jacova C, Robillard A, Garcia A, Chow T, Borrie M et al. Diagnosis and treatment of dementia: 2. Diagnosis. *Can Med Assoc J* 2008;178(7):825-36.
24. Streh D, Mertz M, Knüppel H, Neitzke G, Schmidhuber M. The full spectrum of ethical issues in dementia care: systematic qualitative review. *Br J Psychiatry* 2013;202(6):400-6.
25. Taylor A, Gosney MA. Sexuality in older age: essential considerations for healthcare professionals. *Age Ageing* 2011;40(5):538-43.
26. De Medeiros K, Rosenberg PB, Baker AS, Onyike CU. Improper sexual behaviors in elders with dementia living in residential care. *Dement Geriatr Cogn Disord* 2008;26(4):370-7.
27. Davies J, Gregory D. Entering the dialogue: Marriage biographies and dementia care. *Dementia* 2007;6(4):481-8.
28. Tucker I. Management of inappropriate sexual behaviors in dementia: a literature review. *Int Psychogeriatr* 2010;22(5):683-92.
29. Alagiakrishnan K, Lim D, Brahim A, Wong A, Wood A, Senthilselvan A et al. Sexually inappropriate behavior in demented elderly people. *Postgrad Med J* 2005;81(957):463-6.
30. Stokes G. Challenging behavior in dementia: a person-centered approach. Routledge; 2017.
31. Sadowsky CH, Galvin JE. Guidelines for the management of cognitive and behavioral problems in dementia. *J Am Board Fam Med* 2012;25(3):350-66.
32. Archibald C. Sexuality and dementia in residential care – whose responsibility ?. *Sex Relat Ther* 2002;17(3):301-9.

33. Buhr GT, White HK. Difficult behaviors in long-term care patients with dementia. *J Am Med Dir Assoc* 2007;8(3):e101-13.
34. Ozkan B, Wilkins K, Muralee S, Tampi RR. Pharmacotherapy for inappropriate sexual behaviors in dementia: a systematic review of literature. *Am J Alzheim Dis Dement* 2008;23(4):344-54.
35. Makimoto K, Kang HS, Yamakawa M, Konno R. An integrated literature review on sexuality of elderly nursing home residents with dementia. *Int J Nurs Pract* 2015;21:80-90.
36. Chen ST. Treatment of a patient with dementia and inappropriate sexual behaviors with citalopram. *Alz Dis Assoc Disord* 2010;24(4):402-3.
37. Joller P, Gupta N, Seitz DP, Frank C, Gibson M, Gill SS. Approach to inappropriate sexual behavior in people with dementia. *Can Fam Phys* 2013;59(3):255-60.
38. Srinivasan S, Weinberg AD. Pharmacologic treatment of sexual inappropriateness in long-term care residents with dementia. *Ann Long Term Care* 2006;14(10):20-30.
39. Furlan JC, Henri-Bhargava A, Freedman M. Clomipramine in the treatment of compulsive behavior in frontotemporal dementia: a case series. *Alz Dis Assoc Disord* 2014;28(1):95-8.
40. Manoochehri M, Huey ED. Diagnosis and management of behavioral issues in frontotemporal dementia. *Curr Neurol Neurosci Rep* 2012;12(5):528-36.
41. Teranishi M, Kurita M, Nishino S, Takeyoshi K, Numata Y, Sato T et al. Efficacy and tolerability of risperidone, yokukansan, and fluvoxamine for the treatment of behavioral and psychological symptoms of dementia: a blinded, randomized trial. *J Clin Psychopharmacol* 2013;33(5):600-7.
42. Teranishi M, Kurita M, Nishino S, Takeyoshi K, Numata Y, Sato T et al. Efficacy and tolerability of risperidone, yokukansan, and fluvoxamine for the treatment of behavioral and psychological symptoms of dementia: a blinded, randomized trial. *J Clin Psychopharmacol* 2013;33(5):600-7.
43. Ward RF, Manchip S. 'Inappropriate' sexual behaviors in dementia. *Rev Clin Gerontol* 2013;23(1):75-87.
44. Trifirò G, Sini G, Sturkenboom MC, Vanacore N, Mazzaglia G, Caputi AP et al. Prescribing pattern of antipsychotic drugs in the Italian general population 2000–2005: a focus on elderly with dementia. *Int Clin Psychopharmacol* 2010;25(1):22-8.
45. Cipriani G, Ulivi M, Danti S, Lucetti C, Nuti A. Sexual disinhibition and dementia. *Psychogeriatrics* 2016;16(2):145-53.
46. Bossini L, Casolaro I, Koukouna D, Cecchini F, Fagiolini A. Off-label uses of trazodone: a review. *Exp Opin Pharmacother* 2012;13(12):1707-17.
47. Pike CJ, Carroll JC, Rosario ER, Barron AM. Protective actions of sex steroid hormones in Alzheimer's disease. *Front Neuroendocrinol* 2009;30(2):239-58.
48. Rocca WA, Mielke MM, Vemuri P, Miller VM. Sex and gender differences in the causes of dementia: a narrative review. *Maturitas* 2014;79(2):196-201.
49. Abdo CH. Sexuality and couple intimacy in dementia. *Curr Opin Psychiatry*. 2013;26(6):593-8.
50. Geerlings MI, Stroyk D, Masaki K, Remaley AT, Petrovitch H, Ross GW, et al. Endogenous sex hormones, cognitive decline, and future dementia in old men. *Ann Neurol* 2006;60(3):346-55.
51. Desai AK, Schwartz L, Grossberg GT. Behavioral disturbance in dementia. *Curr Psychiatr Rep* 2012;14(4):298-309.
52. Aarsland D, Sharp S, Ballard C. Psychiatric and behavioral symptoms in Alzheimer's disease and other dementias: etiology and management. *Curr Neurol Neurosci Rep* 2005;5(5):345-54.
53. Freymann N, Michael R, Dodel R, Jessen F. Successful treatment of sexual disinhibition in dementia with carbamazepine. *Pharmacopsychiatry* 2005;38(03):144-5.
54. Raina P, Santaguida P, Ismaila A, Patterson C, Cowan D, Levine M et al. Effectiveness of cholinesterase inhibitors and memantine for treating dementia: evidence review for a clinical practice guideline. *Ann Intern Med* 2008;148(5):379-97.
55. Sonkusare SK, Kaul CL, Ramarao P. Dementia of Alzheimer's disease and other neurodegenerative disorders—memantine, a new hope. *Pharmacol Res* 2005;51(1):1-7.