

# COMPARATIVE ASSESSMENT OF FACIAL ASYMMETRY IN MALOCCLUSION USING POSTEROANTERIOR VIEW

Dr. Anshu Sahu Dr. Rajeev Lall, D N Kapoor, D.L Naidu

**ABSTRACT:**

**Introduction:** The study was conducted to assess facial asymmetry in various dental malocclusions and to determine asymmetry in lower, mid and upper face and jaws using postero-anterior cephalometric analysis. **Method:** 120 postero-anterior cephalograms were taken of individuals between 12 to 25 years age group of both sexes, and were divided into 4 groups, Angle's class I excellent occlusion, Angle's class I malocclusion, Angle's class II malocclusion and Angle's class III malocclusion. These cephalograms were traced and Grummon's analysis was performed. **Results:** In Angle's class I occlusion and Angle's class II malocclusion the results obtained showed asymmetry was present in the upper face. Correlation was found between occlusion, malocclusion and facial asymmetry. **Conclusion:** Facial asymmetry was found in all dental occlusions whether excellent or malocclusion group, with maximum asymmetry having upward trend towards upper face starting from lower.

**KEY WORDS:** Facial Asymmetry, Posteroanterior Cephalogram, Craniofacial Skeleton.

**INTRODUCTION:** Asymmetry in the craniofacial areas can be recognized as differences in the size or relationships of the two sides of the face. This may be the result of discrepancies either in the form of individual bones or a malposition of one or more bones in the craniofacial complex. The asymmetry may also be limited to the overlying soft tissues<sup>1</sup>.

Facial asymmetries are imbalances that occur between the homologous parts of the face affecting the proportion of these parts to one another with regard to size, form and position on opposite sides of the plane, line or point. Facial asymmetry exists in orthodontic as well as non-orthodontic individuals. Because facial asymmetries are very often present with dental asymmetries they are of clinical importance in the treatment of malocclusions of the teeth<sup>2</sup>.

Lundstrom<sup>4</sup> explained that asymmetry




FIG 1: Showing combined asymmetric variables of all four groups.




Fig.2: Showing the image of posteroanterior cephalogram tracing with the landmarks used in the analysis.

Fig. 3: Showing constructed lines used for linear asymmetry assessment MSR (red), and lines Z-MSR, ZA-MSR, NC-MSR, J-MSR, A6-MSR, B6-MSR, AG-MSR (blue) and lines used for mandibular deviation assessment M-MSR (orange).

can be genetic or non-genetic in origin and that it is usually a combination of both. Asymmetries can be classified according to the structures that are involved. Dental asymmetries can be caused by local factors such as early loss of primary teeth, congenitally miss-

ing teeth, and habits such as thumb sucking. Lack of exactness in genetic expression affects the teeth on the right and left sides, causing asymmetries in mesiodistal crown diameters<sup>4</sup>.

The aims and objectives of the study were

# CLINICAL SECTION

to assess the asymmetry in lower, mid and upper face and jaws using posteroanterior cephalometry, and to ascertain the correlations between occlusion and facial asymmetry.

**MATERIALS AND METHOD:** In all, 120 subjects were selected from the population of Moradabad city of Uttar Pradesh, India using the variables as defined below. The sample selected ranged in the age group of 12 to 25 years, in both sexes. Selected individuals were subjected to cephalometric radiography in the department of oral medicine and radiology using a cephalostat of a cephalometric machine manufactured by Villa Systemi (Italy).

The selected subjects had Angle's class I excellent occlusion, Angle's class I malocclusion, Angle's class II malocclusion, Angle's class III malocclusion and were named as group A, B, C and D respectively. These subjects were selected on the following basis-

1. Harmonious & apparently symmetrical face.
2. Full complement of teeth with good posterior interdigitation excluding 3rd molars.
3. No history of trauma.
4. No history of prior orthodontic or surgical treatment.
5. No evidence of TMJ dysfunction or congenital TMJ ankylosis.

The parameters used for group A were detailed as having Angle's Class I molar relation, overjet = 2-4mm, overbite = 2-4mm, symmetrical upper and lower arch, spacing = 0-2mm, contact point displacement = 0-2mm and rotation = mild degree 5. The subject's name, age, sex were recorded and consent was taken, following clinical examination, their posteroanterior cephalogram were taken using standardized technique. The exposure parameters were 75 KVP, 10 mA and exposure time was 1.60 seconds.

Following landmarks and planes were included in the study Fig(I) & Table I.

1. Cg-Cristagalli - A vertically elongated diamond shaped radiopacity appearing between the orbital outline on PA cephalogram. Used to establish a MSR (Mid reference line) 7.

2. Z - Zygomatic suture point-Medial and anterior junction of zygomatic bone with frontal bone. [Right and Left] 8.

3. ZA - Centers of the zygomatic arches [Right and Left] 8.

4. J -Jugal process-Lowest point on the curve of zygomatic bone. Also the point on the jugal process of the maxilla at a crossing with the tuberosity of the maxilla, in the frontal 8.




Fig. 4: Showing the maxillomandibular relationships. MSR line (red), CG-J-MSR triangle on left side (pink), CG-AG-MSR triangle on right side (blue).

Fig. 5: Showing the linear vertical discrepancies, MSR line, vertical discrepancies shown in blocks.

## TABLES

Table No I showing cephalometric line

1) Reference (vertical line)- MSR- mid sagittal reference line. <sup>9</sup>
2) Maxillary width (horizontal line) - J to MSR jugale- crossing of the outline of the tuberosity with the outline of the jugal process. (the medial aspects of the jugal processes). <sup>9</sup>
3) Nasal cavity width (horizontal width)- NC to MSR widest points in nasal capsule. <sup>9</sup>
4) Mandibular width (horizontal line)- Ag to MSR antegonian- trihedral eminence above gonial notch. <sup>9</sup>

Table No II - Showing Mean, SD, SEM, CV and t-test in Group A Having Class-I Excellent Occlusion for Assessing Facial Asymmetry in Different variables.

Horizontal	MEAN	MEAN	SD	SD	SEM	SEM	CV	CV	t-Test
	Left	Right	Left	Right	Left	Right	Left	Right	
Z-MSR	47.87	49.36	2.12	3.02	0.36	0.51	4.43	6.11	-4.51***
ZA-MSR	66.67	67.73	3.39	4.27	0.57	0.72	5.09	6.30	-1.64
NC-MSR	14.97	15.69	1.49	2.04	0.25	0.34	9.98	12.98	-1.66
J-MSR	32.96	33.50	2.22	2.29	0.38	0.39	6.75	6.83	-1.43
A6-MSR	30.43	30.83	2.25	2.54	0.38	0.43	7.39	8.23	-0.98
B6-MSR	30.39	31.00	2.28	2.38	0.39	0.40	7.50	7.69	-1.21
Ag-MSR	43.84	43.27	3.48	3.19	0.59	0.54	7.93	7.37	0.84

Level of Significance of t-test is >0.05 is N.S.; <0.05 is 2.03\*; <0.01 is 2.73\*\* and <0.001 is 3.60\*\*\* at 33 df.

5. NC -Lateral most point on inside surface of bony nasal cavity [Right and Left] 8.

Table No III - Showing Mean, SD, SEM, CV and t-test in Group B Having Class-I Malocclusion for Assessing Facial Asymmetry in Different variables

Horizontal	MEAN	MEAN	SD	SD	SEM	SEM	CV	CV	t-Test
	Left	Right	Left	Right	Left	Right	Left	Right	
Z-MSR	47.77	48.95	3.18	3.03	0.58	0.55	6.65	6.18	-2.39
ZA-MSR	67.63	66.72	4.01	4.12	0.73	0.75	5.93	6.17	1.31
NC-MSR	15.87	15.50	2.19	2.48	0.40	0.45	13.82	16.03	0.77
J-MSR	33.98	33.52	2.25	2.83	0.41	0.52	6.61	8.45	0.89
A6-MSR	30.90	29.20	2.56	5.95	0.47	1.09	8.28	20.39	1.69
B6-MSR	30.62	29.92	2.65	2.93	0.48	0.54	8.66	9.81	1.23
Ag-MSR	43.95	41.73	3.96	3.55	0.72	0.65	9.02	8.51	2.31

Level of Significance oft-test is >0.05 is N.S;<0.05 is 2.05\*; <0.01 is 2.76\*\*and <0.001 is 3.67\*\*\* at 28 df.

Table No. IV Showing Mean, SD, SEM, CV and t-test in Group C Having Class-II Malocclusion for Assessing Facial Asymmetry in Different variables.

Horizontal	MEAN	MEAN	SD	SD	SEM	SEM	CV	CV	t-Test
	Left	Right	Left	Right	Left	Right	Left	Right	
Z-MSR	46.87	48.10	3.57	3.68	0.65	0.67	7.63	7.66	-2.67*
ZA-MSR	65.27	65.93	5.43	5.14	0.99	0.94	8.33	7.80	-0.68
NC-MSR	15.53	16.25	3.46	3.84	0.63	0.70	22.27	23.61	-1.50
J-MSR	31.80	32.32	4.85	5.13	0.89	0.94	15.25	15.86	-1.33
A6-MSR	30.33	30.17	3.45	3.71	0.63	0.68	11.37	12.29	0.27
B6-MSR	29.65	29.73	3.31	3.15	0.60	0.57	11.17	10.59	-0.14
Ag-MSR	42.37	42.52	4.37	3.85	0.80	0.70	10.33	9.05	-0.20

Level of Significance oft-test is >0.05 is N.S;<0.05 is 2.05\*; <0.01 is 2.76\*\*and <0.001 is 3.67\*\*\* at 28 df.

6. Ag-Antegonion. Highest point in the antegonial notch. Antegonial point on the mandibular border at lower margin of trihedral eminence above gonial notch [Right and Left].9

7. A6-Upper first permanent molar. In the frontal [cephalogram] it is the buccal-most point on the crown of upper first molar8.

8. B6-Lower first permanent molar. Frontally it is the buccal most point on the crown of the lower molar8.

9. Me- Menton. Lower most point of the contour of the chin8 .

**Areas for maxillomandibular comparison:**  
Maxillary- Cg - J - MSR.  
Mandibular - Cg - Ag - MSR.

Horizontal asymmetry assessment involved measurement of the horizontal lines which were the perpendicular projections of the bilateral landmarks on the MSR, i.e. Z-MSR, ZA-MSR, NC-MSR, J-MSR, A6-MSR, B6-MSR, and Ag-MSR and were measured for right and left side. A difference in reading of right and left side of a pair of landmarks provided the horizontal asymmetry of the landmarks. Fig (II).

The vertical lines between the points of perpendicular projections on MSR were drawn to depict any vertical discrepancy between the landmarks of right and left side. A vertical difference in the left and right points provided the vertical asymmetry of the landmarks. Fig(IV)

Mandibular deviation was assessed by measuring the linear horizontal distance, between the points of line MSR falling on the lower border of the mandible and Menton. Fig (II)

Four lines were constructed, perpendicular to MSR, from Ag and from J, bilaterally. Lines connecting Cg and J and lines from Cg to Ag were also drawn. Two pairs of triangles are constructed and each pair is bisected by MSR. Their areas were calculated and compared to that of the opposite side. Fig(III)

**RESULTS AND DISCUSSION:**

Table II Group A is showing Mean, CV, SD, SEM for both right and left: side. When mean values of all parameters are compared between right and left side, it is observed that Z to MSR variables are significantly different at 5 % level of significance.

Whereas among other parameter insignificant difference was observed, indicating that in most of the variables there is no evidence of any facial asymmetry.

Table III Group B is showing Mean, CV, SD, SEM for both right and left side.

When mean values of all parameters are compared between right and left side, it is observed that Z to MSR and Ag to MSR variables are significantly different at 5 % level of significance. Whereas among other parameter insignificant difference was observed, indicating that in most of the variables there is no evidence of any facial asymmetry in individual having Angle's Class I malocclusion.

Table IV Group C is showing Mean, CV, SD, SEM for both right and left side. When mean values of all parameters are compared between right and left side, it is observed that Z to MSR variables are significantly different at 5 % level of significance. Whereas among other parameter insignificant difference was observed, indicating that in most of the variables there is no evidence of any facial asymmetry in individual having Angle's Class II malocclusion.

Table V Group D is showing Mean, CV, SD, SEM for both right and left side. When mean values of all parameters are compared between right and left side, it is observed that J to MSR variables are significantly different at 5 % level of significance. Whereas, among other parameters, insignificant difference was observed, indicating that in most of the variables there is no evidence of any facial asymmetry in individual having Angle's Class III malocclusion.

Table VI shows variables having asymmetry in all four groups A, B, C and D.

Table VI and figure V show variable having highly significant difference in Z- MSR between right and left measurement in group

# CLINICAL SECTION

A and same is true in Group B - C whereas insignificant difference exists in Angles class III malocclusion. Measurement Ag-MSR in Group Band J-MSR in Group D both are showing significant difference at 5% which is an indication of asymmetry.

Table VII showing the Mean, CV, SD, SEM between right and left maxillomandibular comparison measurement, using CG-AG-MSR and CG-J-MSR to assess the asymmetry.

From the table, it is observed that measurement CG-J-MSR is showing significant difference at 5% level for right and left side indicating thereby asymmetry in maxillary region for group C and variable CG-AG-MSR in group B.

For rest of the groups, in maxillo-mandibular comparison, measurements are having insignificant difference at 5% level for right and left side indicating thereby symmetry between right and left side. The coefficient of variation was also found to be consistent in all parameters used to assess asymmetry.

Table VIII shows the mean standard deviation, SEM and CV of variables used to assess the asymmetry in all the four groups. The coefficient of variation was found to be variable in a parameter used to assess mandibular deviation. The CV was found to be highest in group B, compared to other groups.

Table IX shows excellent occlusion having molar class I relation correlated for different variables having malocclusion for asymmetry in horizontal, vertical plane and difference in maxillomandibular comparison measurements and mandibular deviations.

There are four types of dentofacial asymmetries studied by Cheney<sup>10</sup> among which the vertical displacements are asymmetrical variations which result from height difference in size shape, and/or form between dentofacial parts on the two sides of the face.

Thompson<sup>11</sup> studied facial symmetry and stated that, it must be recognized that there is no truly symmetrical face regardless of race, age or period of an individual. Fischer<sup>2</sup> noted that the factors responsible for asymmetries in the dentofacial complex are not confined to the teeth and alveolar process. They may be found in the various components parts of the face and all the structures surrounding the teeth.

It was seen that, variable Z-MSR in the group A,B,C; Ag - MSR of class I malocclusion and J- MSR of group D showed significant difference in comparison of right and left side. The right side was more deviated as compared to left side. This was in accordance with the previous study done by Haraguchi S et al<sup>12</sup> in 2002 who stressed that the frequent laterality of face may be ascribed to the

Table No. V Showing Mean, SD, SEM, CV and t-test in Group D Having Class-III malocclusion for assessing facial asymmetry in different variables

HORIZONTAL	MEAN	MEAN	SD	SD	SEM	SEM	CV	CV	t-Test
	Left	Right	Left	Right	Left	Right	Left	Right	
Z-MSR	47.90	48.32	3.21	3.88	0.64	0.78	6.70	8.03	-0.58
ZA-MSR	66.04	66.22	4.68	4.70	0.94	0.94	7.09	7.10	-0.21
NC-MSR	15.04	15.44	1.43	1.77	0.29	0.35	9.50	11.46	-0.95
J-MSR	32.70	33.60	3.15	3.22	0.63	0.64	9.64	9.60	-2.34*
A6-MSR	29.68	30.34	2.33	3.16	0.47	0.63	7.84	10.41	-1.28
B6-MSR	29.84	30.60	2.63	3.11	0.53	0.62	8.81	10.17	-1.43
Ag-MSR	42.76	43.36	4.02	4.30	0.80	0.86	9.41	9.93	-0.58

Level of Significance oft-test is >0.05 is N.S.;<0.05 is 2;07\*; <0.01 is 2.81 \*\*and <0.001 is 3.77\*\*\* at 23 df.

Table No VI Showing variables having asymmetry in all four groups

Group	GroupA	GroupB		Group C	Group D
Variables	Z-MSR	Z-MSR	Ag-MSR	Z-MSR	J-MSR
Difference of mean	1.49	-1.18	2.22	-1.24	-0.90
SD	1.95	2.71	5.26	2.53	1.93
SEM	0.33	0.50	0.96	0.46	0.39
t test	-4.51 **	-2.39*	2.31 *	-2.67*	-2.34*
CV	131.25	-229.16	237.51	-204.86	-213.97

FOR Group A, Level of Significance of t-test is >0.05 is N.S.;<0.05 is 2.03\*; <0.01 is 2.73\*\*and <0.001 is 3.60\*\*\* at 33 df.

FOR Group B & C, Level of Significance of t-test is >0.05 is N.S.;<0.05 is 2.05\*; <0.01 is 2.76\*\*and <0.001 is 3.67\*\*\* at 28 df.

FOR Group D - Level of Significance of t-test is >0.05 is N.S.;<0.05 is 2.07\*; <0.01 is 2.81 \*\*and <0.001 is 3.77\*\*\* at 23 df.

dominant growth potential of the jaw's right side.

It was observed that measurement CG-J-MSR is showing significant difference for right and left side indicating thereby asymmetry in maxillary region for group C. Bjork suggested that there is a slight tendency for most of the cranial bones to be larger on the right side in the underformed (normal) crania. Asymmetry of upper face occurs to prevent midline deviations. This means that at the cost of maintenance of midline, asymmetry of the face results.

Measurement CG-Ag-MSR is showing significant difference between right and left side indicating thereby asymmetry in mandibular region for group B. In accor-

dance with the study done by Haraguchi S et al<sup>12</sup> there was a general tendency of the inferior landmarks to deviate more frequently and at greater distances than the more superiorly located landmark because growth of mandible is largely seen at the condylar region, the mandible is likely to show gradual deviation during growth period, as if it swings with a condylar head on the affected side as its centre of rotation. Chierici et al<sup>16</sup> described this gradual deviation with the help of an animal experiments and stressed asymmetry of the face is related to functional demands of the masticatory apparatus and the musculoskeletal systems. Skeletal asymmetry reflects onto the soft tissue of the face. In this study asymmetry was obvious in the

Table NoVII Showing mean, SD, SEM, CV, and t-test in all groups for variable CG-J-MSR and CG-AG-MSR to assess asymmetry

CG-J-MSR									
Groups.	MEAN		SD		SEM		CV		t-Test
	LEFT	RIGHT	LEFT	RIGHT	LEFT	RIGHT	LEFT	RIGHT	
A	1057.53	1069.44	141.99	136.3	24	23.04	13.43	12.74	-0.97
B	1082.75	1086.98	172.8	179.96	31.55	32.86	15.96	16.56	-0.19
C	978.98	1004.43	216.69	206.27	39.56	37.66	22.13	20.54	-2.18*
D	1020.13	1049.41	120.71	142.46	24.14	28.49	11.83	13.58	-2

CG-AG-MSR									
Groups.	MEAN		SD		SEM		CV		t-Test
	LEFT	RIGHT	LEFT	RIGHT	LEFT	RIGHT	LEFT	RIGHT	
A	2299.06	2279.99	249.56	275.92	42.18	46.64	10.85	12.1	0.57
B	2347.05	2244.3	340.5	286.68	62.17	52.34	14.51	12.77	2.25*
C	2186.17	2212.16	364.53	333.71	66.55	60.93	16.67	15.09	-0.68
D	2227.49	2225.63	346.39	314.05	69.28	62.81	15.55	14.11	0.04

FOR Group A - Level of Significance oft-test is >0.05 is N.S.; <0.05 is 2.02\*; <0.01 is 2.70\*\*and <0.001 is 3.46\*\*\* at 53or58or63 df.

FOR Group B & C - Level of Significance of t-test is >0.05 is N.S.; <0.05 is 2.05\*; <0.01 is 2.76\*\*and <0.001 is 3.67\*\*\* at 28 df.

FOR Group D - Level of Significance of t-test is >0.05 is N.S.; <0.05 is 2.07\*; <0.01 is 2.81 \*\*and <0.001 is 3.77\*\*\* at 23 df.

Table No VIII - Mandibular Deviations

	A	B	C	D
Mean	0.6	0.5	0.37	0.6
Std	1.02	1.6	1.02	1.53
Sern	0.1	0.09	0.07	0.12
CV	170.06	319.48	277.24	255.72
Range min	-1	-3	-1.5	-2
Range max	4	5.5	2.5	2.5

upper jaw but could not extend up to the zygoma.

Maximum coefficient of variation in mandibular deviation was noticed in group C, and minimum deviation was seen in the group B. This shows that although individuals have excellent occlusion still they exhibit asymmetry which has been stressed by Utreja<sup>13</sup> in 1973. Also by Sheats<sup>14</sup> who studied prevalence of orthodontic asymmetries stated in a study that among orthodontic patients, the most common asymmetry trait was mandibular midline deviation from the facial midline. Significant correlation was found between group B and group C, in measurement A6 -MSR which demonstrates that as the malocclusion increases in severity

from group A to group C the value of A6 -MSR increases as well.

When correlation was evaluated in vertical variable it was observed that as malocclusion increases from group A to group C, the Z-MSR value increases and when the value of measurement decreases, the correlation demonstrated that as malocclusion increases from group A to group D, which is evident in the measurement of Z-MSR. Similarly, increase in ZA-MSR was observed with increase in severity of malocclusion from group B to group C. Measurement ZA-MSR, NC-MSR, J-MSR demonstrated that with increase in severity of malocclusion from group A to group B, the value of these variables also decreases. Measurements of variable B6-

MSR demonstrates that with increase in severity of malocclusion from group A to group D, the value decreases. For NC-MSR variable it was seen that with increase in severity of malocclusion from group A to group D the value increases.

When correlation was done in the maxilomandibular parameter it was seen that as the severity of malocclusion increases from group A to group C, the value of Cg-J-MSR also increases. Our findings were contrary to the findings of Thompson<sup>11</sup> (1943) in which he observed insignificant difference between malocclusion and asymmetry while Fischer<sup>2</sup> (1954) reported that facial asymmetry was very often present with malocclusion.

Shah and Joshi<sup>15</sup> reported in their study that significantly more subjects were chewing on the right side than on the left side as a matter of habit and since the force of mastication are transmitted from the teeth to the facial and cranial bones, this may be a factor responsible for the right side being larger than the left.

CONCLUSION:

Following conclusions were drawn:

1. Asymmetry of face are common findings in case of all types of dental malocclusion.
2. In Angle's class I occlusion, Angle's class II malocclusion, the results of parameters obtained show that increased asymmetry is present in the upper face, and the asymmetry increases in magnitude as we approach higher in the craniofacial skeleton.
3. Correlation was found between occlusion, malocclusion and facial asymmetry.

REFERENCES:

1. Edward H. Angle; The Dental Cosmos; Classification of malocclusion, 1899; 248-264.
2. Fischer; Asymmetries of the dentofacial complex; Angle orthod, 1954, 24, 4: 179-192.
3. P.S. Vig and A. B. Hewitt; Asymmetry of the human facial skeleton, Angle Orthod, 1975,45,2: 125-129.
4. Lundstrom, A.; Some asymmetries of dental arches, jaws and skull and their etiological significance, Am J Orthod Dentofac Orthop, 1961,47: 81-106.
5. Little RM; The irregularity index: The quantitative score of mandibular anterior alignment. Am J Orthod Dentofac Orthop, 68, 5: 554-563.
6. White S.c., Pharoah M.J.; Oral Radiology - Principles and interpretation, edition 5th, 2008, Mosby: 210-213.
7. Grummons D. C. and Kappeyne Van De Copello; A frontal asymmetry analysis, J ClinOrthod, 1987,21: 448-65.

## CLINICAL SECTION

8. Kunhiko Miyashita; Contemporary cephalometric radiography; first edition; Quintessence publishers; 1996, 160-209, 259-261.,
9. Athanasiou A. E.; Data and patterns of transverse dentofacial structures of 6 to 15 year old children: A posteroanterior cephalometric study; Am J Orthod Dentofac Orthop, 1992, 101: 465-71.
10. Edward A. Cheney; Dentofacial asymmetries and their clinical significance; Am J Orthod Dentofac Orthop; 1961,47, 11: 814-829.
11. Thompson JR; Asymmetry of face, JADA, 30, 1943: 1859-1871.
12. Haraguchi S et al; Facial asymmetry in subjects with skeletal class III deformity, Angle Orthod, 2002, 72, 1: 28-35.
13. Utreja A. K.; Craniofacial asymmetry and malocclusion- A posteroanterior cephalometric evaluation. Thesis, 1973, Lucknow Univ.
14. Sheats RD et al; Prevalance of orthodontic asymmetries: Seminars in orthodontics, 1998 Sept; 4, 3: 138-45.
15. Shah and Joshi; An assessment of asymmetry in the normal craniofacial complex: Angle Orthod, 1978,48,2: 141-148.
16. Chierici et al; Morphogenetic experiments in facial asymmetry: the nasal cavity. American journal of physical Anthropology, 1973, 38(2): 291-99.

Table No.IX Showing Correlation Coefficient Between Various Variables With Excellent Occlusion And Malocclusion

CORRELATION IN HORIZONTAL PARAMETERS			
EX Occ	MALOCCLUSION		
	Group A vs Group B	Group A vs Group C	Group A vs Group D
Z- MSR	0.076	0.133	0.239
ZA-MSR	0.33	0.211	0.064
NC-MSR	-0.202	-0.108	0.208
J-MSR	-0.068	0.059	0.194
A6-MSR	0.066	0.252*	0.09
B6-MSR	0.008	0.096	-0.196
Ag - MSR	-0.21	-0.03	-0.053
CORRELATION IN VERTICAL PARAMETERS			
Z- MSR	0.048	0.430***	-0.364***
ZA-MSR	-0.271 *	0.390**	-0.017
NC-MSR	-0.357**	0.213	0.308*
J-MSR	-0.288*	0.026	-0.045
A6-MSR	-0.008	-0.107	0.105
B6-MSR	0.049	-0.08	-0.326**
Ag - MSR	-0.122	-0.191	0.008
CORRELATION IN MAXILLOMANDIBULAR VARIABLE			
CG-AG-MSR	0.23	-0.041	0.011
CG-J-MSR	0.369**	-0.03	-0.108
CORRELATION WITH MANDIBULAR DEVIATION			
M-MSR	-0.016	-0.196	0.001

**DR ANSHU SAHU, DR RAJEEV LALL, KAPOOR D.N., NAIDU D.L.,**

1. Reader, Department of Orthodontics, Hazaribag College of Dental Sciences, Hazaribag, Jharkhand

2. Professor and Head, Department of Orthodontics, Hazaribag College of Dental Sciences, Hazaribag, Jharkhand

3. Professor, Kothiwal Dental College & Research Center, Moradabad, U.P.

4. Professor & Head, Department of Orthodontics, KLR Lenora Institute of Dental Sciences, Rajanagram, Rajahmundry, A.P.

## ICD SECTION VI FELLOWS IN NEWS

### 1. Dr. Mahesh Verma

- Padma Shri was conferred upon Master ICD Past President Dr. Mahesh Verma by the Govt. of India.
- He also took over as President of IDA Head Office.

### 2. Dr. U.S. Krishna Nayak

- Appointed as Dean, Faculty of Dental Sciences NITTE University, Mangalore.
- Nominated by the Govt. of Karnataka to the committee to formulate Oral Health Policy of the State.
- Awarded "Best Dentist Award".
- "Outstanding Professor Award" at the national level by the Indian Orthodontic Society which will be awarded during the forthcoming National Conference of the Indian Orthodontic Society at Kolkata.

### 3. Dr. Rajiv K. Chugh

- Elected President Elect, IDA Delhi State Branch.
- Awarded 'Best Clinician 2014' by Heal Talk Journal.

4. Dr. Santosh Ravindran received the best driver award for high altitude driving - Mahindra monastery drive organised by Mahindra adventures and extreme sports organisation. Delhi - Chandigarh - Manali - Jispa - Sarchu - Leh - Kargil - Sirinagar, this was reported in times of India on Oct 4th, 2014.